

Plant		Oat		34(06010)	Primary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Culm length	10 plants, 2 replications	Measurement	cm (integer)		Length from the ground to the base of panicle at the full heading stage
2	Panicle length	10 plants, 2 replications	Measurement	cm (round to the 1st decimal place)		Length from the base to the tip of panicle (excluding awn)
3	Number of panicles	10 plants, 2 replications	Measurement	Number of panicles /plant (round to the 1st decimal place)		Number of panicles per plant at the full heading stage
4	Glume color	2 plots, 2 replications	Observation	1:Light yellow 2:Yellow-Light yellow 3:Yellow 4:Gray-Yellow 5:Gray 6:Gray-Brown 7:Brown 8:Brown-Black 9:Black		Color of glumes at maturity
5	Weight of 1000 grains	100 seeds, 3 replications	Measurement	g (round to the 1st decimal place)		1000 grain weight estimated by sampling 100 clean grains with 3 replications
6	Heading date	2 plots, 2 replications	Observation	date		Date when 50% of productive tillers have headed
7	Culm thickness	10 plants, 2 replications	Measurement	mm (round to the 1st decimal place)		Long diameter of the middle internode of main stem at a height of 10 to 15 cm at heading stage
8	Leaf length	10 plants, 2 replications	Measurement	cm (round to the 1st decimal place)		Length of the first leaf blade below flag leaf
9	Leaf width	10 plants, 2 replications	Measurement	mm (round to the 1st decimal place)		Width of widest part of the first leaf blade below flag leaf
10	Date of maturity	2 plots, 2 replications	Observation	date		Date when 80% of the necks of panicles turned yellow and grains became as hard as wax

Plant		Oat		34(06010)	Primary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Plant length at early stage	10 plants, 2 replications	Measurement	cm (integer)		Length from the ground to the tip of leaf at the early stage before wintering
2	Plant length in spring	10 plants, 2 replications	Measurement	cm (integer)		Length from the ground to the tip of the uppermost leaf of plant in early spring
3	Plant type	2 plots, 2 replications	Observation	1:Erect 2:Nearly erect 3:Semi-erect 4:Slightly semi-erect 5:Intermediate 6:Slightly intermediate 7:Semi-prostrate 8:Nearly prostrate 9:Prostrate		Angle that outer main stems make with the ground
4	Number of tillers	2 plots, 2 replications	Obs.&Measr.	1:Almost none 2:Extremely few 3:Very few 4:Few 5:Intermediate 6:Some 7:Many 8:Very many 9:Most		Number of tillers per unit area at heading time
5	Texture of culm	2 plots, 2 replications	Observation	1:Almost none 2:Extremely few 3:Very few 4:Few 5:Intermediate 6:Some 7:Rough 8:Very rough 9:Extremely rough		Texture of culms estimated by touching at maturity
6	Pubescence on internode	2 plots, 2 replications	Observation	0:None 1:Almost none 2:Extremely few 3:Very few 4:Few 5:Intermediate 6:Some 7:Many 8:Very many 9:Most		Presence and amount of pubescences on the uppermost internode of main stem
7	Leaf color	2 plots, 2 replications	Observation	1:Extremely light 2:Very light 3:Light 4:Slightly light 5:Intermediate 6:Slightly dark 7:Dark 8:Very dark 9:Extremely dark		Greenness of leaf blades at the full tillering stage
8	Waxiness on leaf sheath	2 plots, 2 replications	Observation	0:None 1:Extremely little 2:Very little 3:Little 4:Slightly little 5:Intermediate 6:Some 7:Much 8:Very much 9:Most		Presence and degree of waxiness on the uppermost leaf sheath of main stem at heading stage
9	Pubescence on leaf sheath	2 plots, 2 replications	Observation	0:None 1:Extremely little 2:Very little 3:Little 4:Slightly little 5:Intermediate 6:Some 7:Much 8:Very much 9:Most		Presence and amount of pubescences on leaf sheath
10	Panicle shape	2 plots, 2 replications	Observation	1:Secund and compressed 5:Intermediate 9:Spreading		Shape of panicle
11	Number of spikelets	10 plants, 2 replications	Measurement	1:Almost none 2:Extremely few 3:Very few 4:Few 5:Intermediate 6:Some 7:Many 8:Very many 9:Extremely many		Number of spikelets per panicle at the full heading date

Plant		Oat		34(06010)	Primary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
12	Spikelet density	2 plots, 2 replications	Observation	1:Extremely sparse 2:Very sparse 3:Sparse 4:Slightly sparse 5:Intermediate 6:Slightly dense 7:Dense 8:Very dense 9:Extremely dense		Density of spikelets on panicle
13	Number of awns	2 plots, 2 replications	Observation	0:None 1:Almost none 2:Extremely few 3:Very few 4:Few 5:Intermediate 6:Some 7:Many 8:Very many 9:Most		Presence and number of awns on a spiklet
14	Awn length	2 plots, 2 replications	Observation	1:Extremely short 2:Very short 3:Short 4:Slightly short 5:Intermediate 6:Slightly long 7:Long 8:Very long 9:Extremely long		Length of the awn of the uppermost caryopsis on panicle
15	Grain shape	2 plots, 2 replications	Observation	1:Extremely slender 2:Very slender 3:Slender 4:Slightly slender 5:Intermediate 6:Slightly thick 7:Thick 8:Very thick 9:Extremely thick		Ratio of the thickness to the length of grains harvested
16	Grain weight per liter	2 plots, 3 replications	Measurement	lg/liter (integer)		Weight of clean grains per liter
17	Grain type	2 plots, 2 replications	Observation	1:Hulled 9:Naked		Presence of hull
18	Pubescence length of caryopsis base	2 plots, 2 replications	Observation	1:Extremely short 2:Very short 3:Short 4:Slightly short 5:Intermediate 6:Slightly long 7:Long 8:Very long 9:Extremely long		Length of pubescences on the base of the uppermost caryopsis on the panicle of main stem
19	Threshability	2 plots, 2 replications	Observation	1:Extremely easy 2:Very easy 3:Easy 4:Slightly easy 5:Intermediate 6:Slightly hard 7:Hard 8:Very hard 9:Extremely hard		Ease of detaching mature caryopses from rachis-branches at maturing stage

Plant		Oat		34(06010)	Secondary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit	Remarks	
1	Crown rust resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high	Degree of the resistance based on the infection of Puccinia coronata by artificial inoculation or late sowing	
2	Lodging resistance	10 plants, 2 replications	Obs.&Measr.	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high	Degree of the resistance based on the measurement of culm strength by instrument or by the degree of lodging by dense planting	

Plant		Oat		34(06010)	Secondary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Holo blight resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the infection of <i>Pseudomonas syringae</i> when the infection is apparent
2	Leaf stripe resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the infection of <i>Pyrenophora avenae</i> when the infection is apparent
3	Scab resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the infection of <i>Gibberella zeae</i> when the infection is apparent
4	Stem rust resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the infection of <i>Puccinia graminis</i> when the infection is apparent
5	Loose smut resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the infection of <i>Ustilago avenae</i> when the infection is apparent
6	Pink snow mold resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the infection of <i>Microdochium nivale</i> when the infection is apparent
7	Pythium snow mold resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the infection of <i>Pythium</i> spp. when the infection is apparent
8	Aphid resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the resistance based on the damage by aphids when the damage is apparent
9	Pre-harvest sprouting	10 plants	Obs.&Measr.	1:Extremely high 2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low 9:Extremely low		Ratio of sprouting grains at maturity in the field

Plant		Oat		34(06010)	Secondary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
10	Cold tolerance	10 plants	Obs.&Measr.	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of cold hardiness based on the rate of survival or winter damage just after overwintering or during winter
11	Tolerance to snow cover	10 plants	Obs.&Measr.	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of the tolerance based on the rate of survivals or winter damage of the heavy snow just after overwintering
12	Tolerance to excess moisture	10 plants	Obs.&Measr.	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Degree of tolerance to wet conditions in the field or in the installation
13	Spring habit	10 plants	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Ratio of heading plants when seeded periodically in spring
14	Regrowth	10 plants	Observation	1:Extremely poor 2:Very poor 3:Poor 4:Slightly poor 5:Intermediate 6:Slightly good 7:Good 8:Very good 9:Excellent		Regrowth in the second week after the first harvest

Plant		Oat		34(06010)	Tertiary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Fresh yield of first harvest	2 plots	Measurement	10 kg/a (integer)		Fresh yield estimated from fresh weight including panicles harvested from an area more than 2 square meters at the first harvest (at the dough-ripe stage for whole crop silage)
2	Dry matter ratio of first harvest	2 plots	Measurement	% (round to the 1st decimal place)		Dry matter ratio measured by sampling 1 kg of fresh weight from cut pieces of more than 5 plants and drying at 70 centi degrees for 48 hours at the first harvest
3	Fresh yield of regrowth	2 plots	Measurement	10 kg/a (integer)		Fresh yield of regrowth after the first harvest measured in the same way as that of the first harvest
4	Dry matter of regrowth	2 plots	Measurement	% (round to the 1st decimal place)		Dry matter ratio of regrowth after the first harvest measured in the same way as that of the first harvest
5	Total fresh yield	2 plots	Measurement	kg/a (integer)		Total of fresh yield harvested during the growing season
6	Total dry matter yield	2 plots	Measurement	kg/a (integer)		Total of dry matter yield during the growing season
7	Grain yield	2 plots	Measurement	kg/a (integer)		Grain yield harvested at dough-ripe stage only for whole crop silage

Plant		Oat		34(06010)	Tertiary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Leaf ratio	2 plots	Measurement	% (round to the 1st decimal place)		Ratio of the weight of leaves in 500 g of fresh sample
2	Hull ratio	2 plots	Measurement	% (round to the 1st decimal place)		Ratio of the weight of hull in 10 g of harvested grains
3	Awning	2 plots	Observation	1:Extremely hard 2:Very hard 3:Hard 4:Slightly hard 5:Intermediate 6:Slightly easy 7:Easy 8:Very easy 9:Extremely easy		Ease of deawning at maturity only for hulled oats
4	Hulling	2 plots	Observation	1:Extremely hard 2:Very hard 3:Hard 4:Slightly hard 5:Intermediate 6:Slightly easy 7:Easy 8:Very easy 9:Extremely easy		Ease of detaching glumes at maturity only for naked oats
5	Dry matter digestibility	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of digestible dry matter analyzed by in vivo test or in vitro enzyme method
6	Crude protein	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of crude protein contained a dry matter base by Kjeldahl method or Near Infra-red Analyzer
7	Acid detergent fiber (ADF)	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of ADF content on a dry matter base analyzed by acid detergent-acetone washing
8	Neutral detergent fiber (NDF)	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of NDF content on a dry matter base analyzed by neutral detergent-acetone washing
9	Acid detergent lignin (ADL)	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of ADL content on a dry matter base analyzed by acid detergent-acetone washing
10	Non-structural carbohydrate (NSC)	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of NSC content on a dry matter base analyzed by phenol-sulfuric acid method after ethanol extraction