

Plant		Chinese milk vetch		39(06015)	Primary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Plant type	10 plants, 2 replications	Observation	1:Erect 2:Nearly erect 3:Semi-erect 4:Slightly semi-erect 5:Intermediate 6:Slightly semi-prostrate 7:Semi-prostrate 8:Nearly prostrate 9:Prostrate		Angle that outer main stems make with the ground
2	Plant height	10 plants, 2 replications	Measurement	cm (round to the 1st decimal place)		Plant length from the ground to the top of plant at the full flowering time
3	Stem thickness	10 plants, 2 replications	Obs.&Measr.	1:Extremely slender 2:Very slender 3:Slender 4:Slightly slender 5:Intermediate 6:Slightly thick 7:Thick 8:Very thick 9:Extremely thick		Diameter of stems at the full flowering time
4	Leaf length	10 plants, 2 replications	Measurement	cm (round to the 1st decimal place)		Length of the biggest leaf from the base of petiole to the tip of top leaflet at the full flowering time
5	Leaflet size	10 plants, 2 replications	Obs.&Measr.	1:Extremely small 2:Very small 3:Small 4:Slightly small 5:Intermediate 6:Slightly large 7:Large 8:Very large 9:Extremely large		Extremely small: $\leq 0.4$ square meter, small: $\leq 0.8$ square meter, intermediate: $\leq 1.2$ square meter, large: $\leq 1.6$ square meter, extremely large: $\geq 2.0$ square meter
6	First flowering date	10 plants, 2 replications	Observation	date		Date when plants began to flower
7	Flower color	10 plants, 2 replications	Observation	1:White 2:Yellowish white 4:Blight red purple 5:Red purple 7:Deep red purple 9:Other		Color of standard and keel petals just after flowering

Plant		Chinese milk vetch		39(06015)	Primary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Stem color	10 plants, 2 replications	Observation	1:Green 3:Light red 5:Red brown 7:Dark red purple 9:Other		Degree of anthocyan pigmentation of stems on the sunny side at flowering time
2	Number of stems	10 plants, 2 replications	Obs.&Mear.	1:Almost none 2:Extremely few 3:Very few 4:Few 5:Intermediate 6:Some 7:Many 8:Very many 9:Abundant		Number of primary branches at the beginning of flowering stage
3	Flower stalk length	10 plants, 2 replications	Measurement	mm (integer)		Average length of 2 flower stalks from 2 longest stems
4	Number of florets per cluster	10 plants, 2 replications	Obs.&Mear.	1:Almost none 2:Extremely few 3:Very few 4:Few 5:Intermediate 6:Some 7:Many 8:Very many 9:Abundant		Number of florets per cluster
5	Pod color	10 plants, 2 replications	Observation	3:Brown 5:Dark brown 7:Black		Color of mature pods
6	Seed shape	10 plants, 2 replications	Observation	1:Round 2:Round-Oval 3:Oval 4:Oval-Elliptic 5:Elliptic 6:Elliptic-Rhombic 7:Rhombic 9:Other		Shape of seeds taken out of mature pods
7	Weight of 1000 seeds	10 plants, 2 replications	Measurement	g / 1000 seeds (round to the 2nd decimal place)		Weight of 1000 seeds estimated by sampling 100 seeds from a mixture of 20 plants, with at least 4 replications

Plant		Chinese milk vetch		39(06015)	Secondary essential character
No	Characters	No. of samples	Methods	Rank or measurement unit	Remarks
1	Sclerotinia wilt rot resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high	Resistance to Sclerotinia trifoliorum based on the degree of damage and the ratio of dead plants when the infection became apparent
2	Overwintering ability	10 plants, 2 replications	Obs.&Measr.	1:Extremely poor 2:Very poor 3:Poor 4:Slightly poor 5:Intermediate 6:Slightly good 7:Good 8:Very good 9:Excellent	Overwintering ability estimated from the rate of survival after overwintering. Extremely poor:less than 10%, poor:less than 30%, intermediate:less than 60%, good:up to 80%, excellent:at least 95%
3	Plant vigor in spring	10 plants, 2 replications	Observation	1:Extremely poor 2:Very poor 3:Poor 4:Slightly poor 5:Intermediate 6:Slightly good 7:Good 8:Very good 9:Excellent	Plant vigor one month after sprouting in spring

Plant		Chinese milk vetch		39(06015)	Secondary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Disease resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Resistance to diseases based on the degree of damage and the ratio of dead plants when the infection became apparent (Note the name of disease)
2	Insect resistance	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Resistance to insects, based on the degree of damage and the ratio of dead plants when the damage became apparent (Note the name of insect)
3	Spring habit	10 plants, 2 replications	Observation	1:Extremely low 2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high 9:Extremely high		Ratio of flowering plants when sown in spring
4	Plant vigor in autumn	10 plants, 2 replications	Observation	1:Extremely poor 2:Very poor 3:Poor 4:Slightly poor 5:Intermediate 6:Slightly good 7:Good 8:Very good 9:Excellent		Amount of growth in late fall

Plant		Chinese milk vetch		39(06015)	Tertiary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Dry matter yield	2 plots	Measurement	kg/a (integer)		Dry matter yield calculated by fresh yield x dry matter ratio/100
2	Dry matter ratio	2 plots	Measurement	% (round to the 1st decimal place)		Ratio of dry matter by sampling 300 g of fresh weight and drying at 70 centi degrees for 48 hours
3	Green yield	2 plots	Measurement	kg/a (integer)		Fresh yield estimated from fresh weight harvested from an area more than 2 square meters
4	Dry matter digestibility	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of digestible dry matter measured by in vivo test or in vitro enzyme method
5	Crude protein content	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of crude protein content on a dry matter base analyzed by Kjeldahl method or Near Infra-red Analyzer
6	Acid detergent fiber content (ADF)	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of ADF content on a dry matter base analyzed by acid detergent-acetone washing
7	Neutral detergent fiber content (NDF)	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of NDF content on a dry matter base analyzed by neutral detergent-acetone washing
8	Acid detergent lignin content (ADL)	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Ratio of ADL content on a dry matter base analyzed by acid detergent-acetone washing
9	Mono-and oligosaccharide content	2 plots, 3 replications	Measurement	% (round to the 1st decimal place)		Mono-and oligosaccharide content on a dry matter base analyzed by thin layer chromatography after ethanol extraction
10	Seed productivity	2 plots, 2 replications	Measurement	g/square meter (round to the 1st decimal place)		Yield of pure seeds per square meter
11	Seed weight per flower head	2 plots, 2 replications	Measurement	mg/flower head (integer)		Weight of pure seeds per flower head, by sampling 20 mature flower heads

Plant	Chinese milk vetch		39(06015)	Tertiary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit	Remarks
12	Seed fertility	2 plots, 2 replications	Measurement	% (round to the 1st decimal place)	Ratio of pure seeds estimated by sampling seeds of 20 mature flower heads