

Plant		Barley		429	Primary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Plant habit	Block	Observation	2:Erect 3:Nearly erect 4:Semi-erect 5:Intermediate 6:Semi-prostrate 7:Nearly-prostrate 8:Prostrate		Plant type before the initiation of internode elongation (January to February). In the district of long snow cover, growth habit is observed before snow cover.
2	Plant type	Block	Observation	2:Normal 8:Uzu		Classified by length and tip shape of coleoptile
3	Culm length	10 plants	Measurement	cm (integer)		Length from ground level to the ear neck of the longest culm
4	Ear length	10 plants	Measurement	cm (round to the 1st decimal place)		Length from ear neck to the top of ear of the longest culm
5	Kernel rows	Block	Observation	2:Two-rowed 5:Coarse six-rowed 8:Six-rowed		Rows of spikelets
6	Existence of awn	Block	Observation	0:Awnless 2:Very scarce 3:Scarce 4:Slightly scarce 5:Intermediate 6:Slightly abundant 7:Abundant 8:Very abundant		Scarce=10%, intermediate=25%, abundant=40%
7	Glume color	Block	Observation	1:Light yellow 2:Yellow 3:Yellowish brown 4:Brown 5:Brownish red 6:Red 7:Reddish purple 8:Purple 9:Deep purple - Black		Glume color at maturity
8	Grain size	Block	Observation	2:Very small 3:Small 4:Slightly small 5:Intermediate 6:Slightly large 7:Large 8:Very large		Grade of grain size
9	Grain color	Block	Observation	0:White 1:Light yellow 2:Yellow 3:Yellow brown 4:Brown 5:Reddish brown 6:Red 7:Reddish purple 8:Purple 9:Dark purple		Color of grain for naked barley
10	Covered or naked kernels	Block	Observation	2:Covered kernels 8:Naked kernels		Classified by detachability of glume
11	Heading date	Block	Observation	date		Date when 40-50% ears have emerged
12	Maturity date	Block	Observation	date		Date when ear necks more than 80% of all ears turn yellow and grains become as hard as wax

Plant	Barley		429	Primary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit	Remarks
13	1000 grain weight	3 replications	Measurement	2:Very light 3:Light 4:Slightly light 5:Intermediate 6:Slightly heavy 7:Heavy 8:Very heavy	Conversion to 1000 grain weight after counting of grains with 20 gram samples (grain moisture: 12.5%)

Plant		Barley		429	Primary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Culm angle	Block	Observation	2:Very close 3:Close 4:Slightly close 5:Intermediate 6:Slightly open 7:Open 8:Very open		Culm angle at the full heading time to maturity. Open type shows a large angle
2	Culm thickness	Block	Observation	2:Very thin 3:Thin 4:Slightly thin 5:Intermediate 6:Slightly thick 7:Thick 8:Very thick		Thickness of culm at maturity
3	Culm stiffness	Block	Observation	2:Very stiff 3:Stiff 4:Slightly stiff 5:Intermediate 6:Slightly tender 7:Tender 8:Very tender		Stiffness of culm at maturity
4	Culm waxiness	Block	Observation	0:Absent 2:Very scarce 3:Scarce 4:Slightly scarce 5:Intermediate 6:Slightly abundant 7:Abundant 8:Very abundant		Degree of culm waxiness in the upper first internode at heading time
5	Presence of auricle	Block	Observation	0:Absent 9:Present		
6	Leaf color	Block	Obs.&Measr.	2:Very light 3:Light green 4:Slightly light 5:Green 6:Slightly dark 7:Dark green 8:Very dark		Leaf color at tillering stage and booting stage or at observation time of growth habit
7	Leaf sheath waxiness	Block	Observation	0:Absent 2:Very scarce 3:Scarce 4:Slightly scarce 5:Intermediate 6:Slightly abundant 7:Abundant 8:Very abundant		Degree of waxiness on the upper first leaf sheath at heading time
8	Leaf sheath pubescence	Block	Observation	0:Absent 2:Very scarce 3:Scarce 4:Slightly scarce 5:Intermediate 6:Slightly abundant 7:Abundant 8:Very abundant		Degree of leaf sheath pubescence
9	Spike configuration	Block	Observation	2:Parallel 3:Slightly parallel 4:Slightly parallel-Intermediate 5:Intermediate 6:Intermediate-somewhat pyramidal 7:Slightly pyramidal-Pyramidal 8:Pyramidal		
10	Spikelet density	10 spikes	Obs.&Measr.	2:Very sparse 3:Sparse 4:Slightly sparse 5:Intermediate 6:Slightly dense 7:Dense 8:Very dense		Number of internodes in rachis (=number of total spikelets - 1)/rachis length (cm)
11	Ear protrusion	5 spikes	Measurement	2:Very short 3:Short 4:Slightly short 5:Intermediate 6:Slightly long 7:Long 8:Very long		Length of the top of flag leaf sheath to spike neck at maturity

Plant		Barley		429	Primary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
12	Spike blade angle	Block	Observation	2:Erect 3:Nearly erect 4:Semi-erect 5:Intermediate 6:Semi-drooping 7:Nearly-drooping 8:Drooping		Degree of spike drooping at maturity
13	Awn length	10 spikes	Obs.&Measr.	2:Very short 3:Short 4:Slightly short 5:Intermediate 6:Slightly long 7:Long 8:Very long		Measurement of the longest 10 awns. Classification compared with standard cultivars
14	Roughness of awn	Block	Observation	2:Very smooth 3:Smooth 4:Slightly smooth 5:Intermediate 6:Slightly rough 7:Rough 8:Very rough		Degree of awn smoothness
15	Hooded awn type	Block	Observation	2:Short hooded awn 5:Hooded awn 8:Long hooded awn		Classified by length of hooded awn
16	Grain shape	Block	Obs.&Measr.	2:Very round 3:Round 4:Slightly oval 5:Oval 6:Slightly slender 7:Slender 8:Very slender		Evaluation based on the ratio of length and width of grain
17	Fineness of husk	Block	Observation	2:Very thin 3:Thin 4:Slightly thin 5:Intermediate 6:Slightly thick 7:Thick 8:Very thick		Judgment by the degree of wrinkling on husk. A wrinkly husk=thin
18	Rachilla hair length	Block	Observation	2:Short 5:Intermediate 8:Long		Observation for malting barley
19	Transverse crease in the lemma base	Block	Observation	0:Absent 9:Present		Presence of transverse crease in the lemma base of grain for malting barley
20	Width of ventral crease	Block	Observation	2:Very narrow 3:Narrow 4:Slightly narrow 5:Intermediate 6:Slightly wide 7:Wide 8:Very wide		Observation for malting barley
21	Lodicule hair length	Block	Observation	2:Short 8:Long		Observation for malting barley
22	Intensity of anthocyanin coloration of flag leaf auricles	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly strong 7:Strong 8:Very strong		Intensity of anthocyanin coloration of flag leaf auricles
23	Frequency of plants with recurved flag leaves	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly high 7:High 8:Very high		Frequency of plants with recurved flag leaves

Plant		Barley		429	Primary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
24	Glaucosity of flag leaf sheath	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly high 7:High 8:Very high		Glaucosity of flag leaf sheath
25	Intensity of anthocyanin of awn tips	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly strong 7:Strong 8:Very strong		Intensity of anthocyanin of awn tips
26	Glaucosity of ear	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly strong 7:Strong 8:Very strong		Glaucosity of ear at flowering to milky stage
27	Attitude of ear	Block	Observation	2:Erect 3:Nearly erect 4:Semi-erect 5:Intermediate 6:Semi-drooping 7:Nearly drooping 8:Drooping		Attitude of ear at milky stage
28	Length of first segment rachis	Block	Measurement	2:Short 5:Intermediate 8:Long		Length of first segment rachis at maturity stage
29	Curvature of first segment rachis	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly strong 7:Strong 8:Very strong		Curvature of first segment rachis at maturity stage
30	Development of sterile spikelets	Block	Observation	2:Non 5:Rudimentary 8:Full		Development of sterile spikelets at maturity stage
31	Attitude of sterile spikelet	Block	Observation	2:Parallel 5:Weakly divergent 8:Divergent		Attitude of sterile spikelet in mid-third of ear at maturity stage
32	Length of glume and its awn relative to grain	Block	Observation	2:Shorter 5:Equal 8:Longer		Length of glume and its awn of median spikelet relative to grain at maturity stage
33	Rachilla hair type	Block	Observation	2:Short 8:Long		Rachilla hair type
34	Anthocyanin coloration of nerves of lemma	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly strong 7:Strong 8:Very strong		Anthocyanin coloration of nerves of lemma at dough ripe stage

Plant		Barley		429	Primary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
35	Spiculation on inner lateral nerves of dorsal side on lemma	Block	Observation	0:Absent 2:Very weak 3:Weak 4:Slightly weak 5:Intermediate 6:Slightly high 7:High 8:Very high		Spiculation on inner lateral nerves of dorsal side on lemma at maturity stage
36	Hairiness of ventral furrow	Block	Observation	0:Absent 8:Present		Hairiness of ventral furrow at maturity stage
37	Disposition of lodicules	Block	Observation	0:Frontal 8:Clasping		Disposition of lodicules at maturity stage
38	Color of aleurone layer	Block	Observation	2:Whitish 5:Weakly colored 8:Strongly colored		Color of aleurone layer at dough ripe stage

Plant		Barley		429	Secondary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Grade of spring habit	10 plants	Obs.&Measr.	1:1 2:2 3:3 4:4 5:5 6:6 7:7		Observation of heading performance after seeds were sown at a constant interval (usually days) from February to April. Accession with high spring habit is classified as 1, with high winter habit is classified as 7.
2	Glutinous or non-glutinous endosperm	Block	Observation	2:Non-glutinous 8:Glutinous		Distinction of glutinous or non-glutinous endosperm based on reaction potassium iodide
3	Awn hulling	Block	Observation	2:Very difficult 3:Difficult 4:Slightly difficult 5:Intermediate 6:Slightly easy 7:Easy 8:Very easy		Investigation at maturity for covered barley
4	Hulling	Block	Observation	2:Very difficult 3:Difficult 4:Slightly difficult 5:Intermediate 6:Slightly easy 7:Easy 8:Very easy		Investigation at maturity for covered barley
5	Threshability	Block	Observation	2:Very difficult 3:Difficult 4:Slightly difficult 5:Intermediate 6:Slightly easy 7:Easy 8:Very easy		Investigation at maturity
6	Lodging resistance	Block	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Synthetic judgment based on the stage of lodging occurrence and degree of lodging
7	Yellow mosaic resistance	10 plants, 2 replications	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Judgment by disease symptom around internode elongation stage and uniformity of heading (middle and southern parts of Japan)
8	Scab resistance	30 plants, 4 replications	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Judgment by the degree of disease infection at dough ripe stage to maturity
9	Powdery mildew resistance	50 plants, 2 replications	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Judgment by the degree and spread of disease symptoms at ripening stage

Plant		Barley		429	Secondary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
1	Leaf rust resistance	Block	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Judgment by the degree and extension of disease symptom at ripening stage
2	Leaf blotch resistance	Block	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Judgment by the degree and extension of disease symptom at ripening stage
3	Time of internode elongation	Block	Observation	2:Very early 3:Early 4:Slightly early 5:Intermediate 6:Slightly late 7:Late 8:Very late		Observation of internode elongation at the end of January to the beginning of April (middle and southern part of Japan)
4	Sprouting resistance	5 spikes	Measurement	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Sprouting of matured spikes under wet condition
5	Cold tolerance	100 plants, 2 replications	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Judgment by the rate of winter-killing and the degree of damage after overwintering (northern part of Japan)
6	Snow mold tolerance	Block	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Judgment by the degree of plant damage after snow melting (snow falling area)
7	Tolerance to moisture	Block	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Endurance under wet condition (note observation stage)
8	Tolerance to soil upheaval	40 plants, 4 replications	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Tolerance to upheaval against frozen soil. Synthetic judgment by the rate of survived plants at two investigation times
9	Resistance to insect pests	Block	Observation	2:Very high 3:High 4:Slightly high 5:Intermediate 6:Slightly low 7:Low 8:Very low		Note insect name

Plant		Barley		429	Tertiary essential character
No	Characters	No. of samples	Methods	Rank or measurement unit	Remarks
1	Yield	Block	Obs.&Measr.	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high	Comparison weight of whole-grains per area with standard cultivars
2	Test weight	3 replications	Measurement	2:Very light 3:Light 4:Slightly light 5:Intermediate 6:Slightly heavy 7:Heavy 8:Very heavy	Weight of 1 liter (grain moisture: 12.5%). Measure 3 times using liter weight vessel.
3	Grain quality	Block	Observation	1:Excellent 2:Very good 3:Good 4:Slightly good 5:Intermediate 6:Slightly poor 7:Poor 8:Very poor 9:Extremely poor	Synthetic judgment of appearance of grain based on fullness, uniformity of size and shape, bright color of grains
4	Glassiness of grain	Block	Obs.&Measr.	3:Powdery 4:Slightly powdery 5:Intermediate 6:Slightly glassy 7:Glassy	Measurement of percentage of glassy kernel. Powdery:<=30% glassy, intermediate:30-70% glassy, glassy:>=70% glassy
5	Whiteness of grain	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high	Measurement of grains with a colormeter, grains are put through 1.8 mm sieve
6	Pearled grain yield	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high	Grains are pearled with a grinder, weight ratio of pearled grain to grain (except for malting barley)
7	Whiteness of pearled grain	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high	Measurement of pearled grains with a colormeter (except for malting barley)
8	Plump-grain percentage	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high	200 gram samples are passed through 2.5 mm sieve for 5 minutes. Weight ratio is measured.
9	Steeping time	Block	Measurement	2:Very short 3:Short 4:Slightly short 5:Intermediate 6:Slightly long 7:Long 8:Very long	Reaching time to 43% grain moisture under water (for malting barley)
10	Malt yield	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high	Weight ratio % of malt to grain in dry weight (for malting barley)

Plant		Barley		429	Tertiary essential character	
No	Characters	No. of samples	Methods	Rank or measurement unit		Remarks
11	Malt extract	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		Weight ratio % of soluble extract to malt (malting barley), measurement by specific gravity of wort
12	Available extract	Block	Calculation	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		Malt extract x Malt yield (for malting barley)
13	Protein content of malt	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		Percentage of total nitrogen content in malt x 6.25 (for malting barley)
14	Soluble nitrogen content of wort	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		Measurement for malting barley
15	Kolbach index	Block	Calculation	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		(Soluble nitrogen content of wort/Total nitrogen content of malt) x 100 (for malting barley)
16	Diastatic power	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		Measurement by starch reaction with enzyme extracted from malt (for malting barley)
17	Diastatic power per total nitrogen	Block	Calculation	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		Diastatic power/Total nitrogen of malt (for malting barley)
18	Apparent final attenuation	Block	Measurement	2:Very low 3:Low 4:Slightly low 5:Intermediate 6:Slightly high 7:High 8:Very high		Calculation based on fermentable carbohydrate of malt extract fermented by yeast in wort
19	Wort color	Block	Measurement	2:Very light 3:Light 4:Slightly light 5:Intermediate 6:Slightly dark 7:Dark 8:Very dark		Measurement of wort color with EBC color desk (for malting barley)

Plant	Barley		429	Tertiary optional character	
No	Characters	No. of samples	Methods	Rank or measurement unit	Remarks
1	Grain hardness	Block	Measurement	2:Very soft 3:Soft 4:Slightly soft 5:Intermediate 6:Slightly hard 7:Hard 8:Very hard	Measurement of grain hardness using Single Kernel Characterization System (SKCS)